

① Problem: IN CLASS IT WAS STATED THAT IF:

25 Points

$$\chi \equiv \frac{x^2}{v_x^2 + s} + \frac{y^2}{v_y^2 + s}$$

AND IF $\rho(x, y) = \hat{\rho}\left(\frac{x^2}{v_x^2 + s} + \frac{y^2}{v_y^2 + s}\right) = \hat{\rho}(\chi) \Big|_{s=0} \equiv \frac{d\eta(\chi)}{d\chi} \Big|_{s=0}$

AND IF $\Psi(x, y) = \frac{-v_x v_y}{4\epsilon_0} \int_0^\infty \frac{\eta(\chi) ds}{\sqrt{v_x^2 + s} \sqrt{v_y^2 + s}}$

THEN IT FOLLOWS THAT $\frac{\partial^2 \Psi}{\partial x^2} + \frac{\partial^2 \Psi}{\partial y^2} = -\frac{\rho(x, y)}{\epsilon_0}$

and so $\Psi(x, y)$ is a solution of Poisson's equation.

a). Calculate $\frac{d\chi}{ds}$. (Note that $x, y, v_x,$ and v_y are held fixed).

b). SHOW THAT $\nabla^2 \Psi = \frac{-v_x v_y}{4\epsilon_0} \int_0^\infty \frac{-4\eta''(\chi) \frac{d\chi}{ds} + 2\eta' \left[\frac{1}{v_x^2 + s} \right] + 2\eta' \left[\frac{1}{v_y^2 + s} \right]}{\sqrt{v_x^2 + s} \sqrt{v_y^2 + s}} ds$

c). Integrate the first ^{term} in the integral ABOVE BY PARTS AND SHOW THAT ~~WHOLE~~ PART OF IT CANCELS THE TERM PROPORTIONAL TO η' .

d). EVALUATE THE INTEGRATED TERM TO SHOW THAT

$$\nabla^2 \Psi = -\frac{\rho}{\epsilon_0}$$

25 Points

(2) Plot $\log(\lambda_{max})$ vs. $\log(\text{ion energy})$ for a heavy ion beam ($m_{mass} = 200$) between 10 keV and 1 GeV. (λ_{max} is the maximum transportable line charge density assuming negligible emittance. Ion energy $\equiv qV$.) Plot for each of the following focusing devices:

- Solenoids
- Electric Quads
- Magnetic Quads
- Einzel lenses

Assume $B_{solenoid} = B_{quad} = 2T$

$\Phi_{lens} = \Phi_{quad} = \pm 100 \text{ kV}$

$v_{beam} \leq 6 \text{ cm}$


Lenz $\gg v_{beam}$

Occupancy $\eta = 0.7$

$$\frac{v_{beam}}{v_{ire}} = 0.7$$

(Use formulas given in class notes on FINAL PAGE OR JOHN'S 6/14/2011 LECTURE).

8/ Consider a Solenoidal Periodic Lattice



- L_p = lattice period
- ηL_p = length solenoids
- $(1-\eta)L_p$ = length drift.
- \hat{R} = solenoid strength
- η = solenoid occupancy $0 < \eta \leq 1$

Larmor Frame

a) Write n transfer matrices $\bar{M}(s|s_i)$ for each section of the periodic lattice in terms of $\Theta = \sqrt{\hat{R}} \eta L_p / 2$ and η .

- \bar{M}_F : Transfer through Solenoid
- \bar{M}_D : " " " Drift

Larmor Frame

b) Write the n transfer matrix $\bar{M}(s_i+L_p, s_i)$ through one lattice period starting from the Solenoid.

Larmor Frame

c) Show that the n phase advance δ_0 of a particle through the lattice period:

$$\cos \delta_0 = \frac{1}{2} \text{Trace } M(s_i+L_p|s_i)$$

can be expressed as:

$$\cos \delta_0 = \cos(2\Theta) - \frac{1-\eta}{\eta} \Theta \sin(2\Theta)$$

$$\Theta = \sqrt{\hat{R}} \eta L_p / 2$$

TPD Problem 8

S.M. Lund P8a/

d) Will it matter where the lattice period is started in the calculation for δ_0 in part c)? Why?

e) For $\theta \ll 1$ (thin lens limit) show that

$$\cos \delta_0 \approx 1 - \frac{\eta |\hat{R}| L_p^2}{2}$$

f) If $\delta_0 \ll 1$, and $\eta \ll 1$, show that

$$\delta_0 \approx \sqrt{\eta |\hat{R}|} L_p$$

g) If one wanted to model a solenoidal focusing lattice by a continuous focusing channel with $R(s) = k_{po}^2 = \text{const}$, how could one choose k_{po}^2 based on part f)?

TPD Problem 9 Problem 4, 25 Points S.M. Lund

P9/

9/ In class we derived the single-particle Courant-Snyder invariant:

$$\gamma x^2 + 2\alpha x x' + \beta x'^2 = \epsilon = \text{const.}$$

where:

$$\beta(s) = W^2(s)$$

$$\alpha(s) = -W(s)W'(s)$$

$$\gamma(s) = \frac{1}{W^2(s)} + W'(s)^2 = \frac{1 + \alpha^2(s)}{\beta(s)}$$

Derive the critical values of the ellipse indicated on the figure below:




Fig. 5.22. Phase space ellipse

From Wiedemann

152

Hint: to avoid messy algebra, take a differential of the constraint equation $\gamma x^2 + 2\alpha x x' + \beta x'^2 = \text{const}$ and use this result to find turning points.

$$\Rightarrow 2\gamma x dx + 2\alpha x dx' + 2\alpha x' dx + 2\beta x' dx' = 0$$

These results are important in understanding the kv distribution derived later to model beams with space-charge

