John Barnard
Steven Lund
USPAS
June 2006

II. Envelope Equations

Paraxial Ray Equation

Envelope equations for axially symmetric beams

Cartesian equation of motion

Envelope equations for elliptically symmetric beams

START WITH NEWTON'S EQUATION WITH THE LORENTZ FORCE:

$$\frac{d\underline{p}}{dt} = q(\underline{E} + \underline{v} \times \underline{B})$$

In cartesian coordinates:

$$\frac{d}{dt}(\gamma m \dot{x}) = \dot{\gamma} m \dot{x} + \gamma m \ddot{x} = q(E_x + \dot{y} B_z - \dot{z} B_y)$$

$$\frac{d}{dt}(\gamma m \dot{y}) = \dot{\gamma} m \dot{y} + \gamma m \ddot{y} = q(E_y + \dot{z} B_x - \dot{x} B_z)$$

$$\frac{d}{dt}(\gamma m \dot{z}) = \dot{\gamma} m \dot{z} + \gamma m \ddot{z} = q(E_z + \dot{x} B_y - \dot{y} B_x)$$

In cylindrical coordinates: (use $\frac{d\hat{e}_r}{dt} = \hat{e}_\theta \dot{\theta}$; $\frac{d\hat{e}_\theta}{dt} = -\hat{e}_r \dot{\theta}$)

$$\frac{d}{dt}(\gamma m \dot{r}) - \gamma m r \dot{\theta}^2 = q(E_r + r\dot{\theta} B_z - \dot{z} B_\theta) \qquad \text{(I)}$$

$$\frac{1}{r}\frac{d}{dt}(\gamma m r^2 \dot{\theta}) = q(E_\theta + \dot{z} B_r - \dot{r} B_z) \qquad \text{(II)}$$

$$\frac{d}{dt}(\gamma m \dot{z}) = q(E_z + \dot{r} B_\theta - r\dot{\theta} B_r) \qquad \text{(III)}$$

<u>When $\frac{\partial}{\partial \theta} = 0$:</u>

$$\underline{E} = -\nabla\Phi - \frac{\partial \underline{A}}{\partial t} = \hat{e}_r\left[-\frac{\partial \Phi}{\partial r} - \frac{\partial A_r}{\partial t}\right] + \hat{e}_\theta\left[-\frac{\partial A_\theta}{\partial t}\right] + \hat{e}_z\left[-\frac{\partial \Phi}{\partial z} - \frac{\partial A_z}{\partial t}\right]$$

$$\underline{B} = \nabla \times \underline{A} = \hat{e}_r\left[-\frac{\partial}{\partial z}(A_\theta)\right] + \hat{e}_\theta\left[\frac{\partial A_r}{\partial z} - \frac{\partial A_z}{\partial r}\right] + \hat{e}_z\left[\frac{1}{r}\frac{\partial}{\partial r}(r A_\theta)\right]$$

$$q r(E_\theta + \dot{z} B_r - \dot{r} B_z) = q\left(-\frac{\partial r A_\theta}{\partial t} - \dot{z}\frac{\partial r A_\theta}{\partial z} - \dot{r}\frac{\partial}{\partial r}(r A_\theta)\right)$$

$$= -q\left[\frac{\partial}{\partial t} + \underline{v}\cdot\frac{\partial}{\partial \underline{x}}\right](r A_\theta)$$

$$= -q\frac{d(r A_\theta)}{dt} \qquad \text{(IV)}$$

Eqtn II & IV $\Rightarrow$ $\frac{d}{dt}(\gamma m r^2 \dot{\theta} + q r A_\theta) = 0$

$$\underline{p} = p_r \hat{e}_r + p_\theta^* \hat{e}_\theta + p_z \hat{e}_z$$

where $p_r = \gamma m \dot{r}$

$p_\theta^* = \gamma m r \dot{\theta}$

$p_z = \gamma m \dot{z}$

$$\frac{d\underline{p}}{dt} = \dot{p}_r \hat{e}_r + p_r \dot{\hat{e}}_r + \dot{p}_\theta^* \hat{e}_\theta + p_\theta^* \dot{\hat{e}}_\theta + \dot{p}_z \hat{e}_z$$

$$\Rightarrow \frac{d\underline{p}}{dt} = (\dot{p}_r - p_\theta^* \dot{\theta}) \hat{e}_r + (p_r \dot{\theta} + \dot{p}_\theta^*) \hat{e}_\theta + \dot{p}_z \hat{e}_z$$

WHERE WE HAVE USED:

$$\frac{d\hat{e}_r}{dt} = \hat{e}_\theta \dot{\theta} \quad \& \quad \frac{d\hat{e}_\theta}{dt} = -\hat{e}_r \dot{\theta}$$

$$\Rightarrow \frac{d\underline{p}}{dt} = \left[\frac{d}{dt}(\gamma m \dot{r}) - \frac{d}{dt}(\gamma m r \dot{\theta}^2)\right] \hat{e}_r$$

$$+ \underbrace{\left[\gamma m r \dot{\theta} + \frac{d}{dt}(\gamma m r \dot{\theta})\right]}_{= \frac{1}{r}\frac{d}{dt}(\gamma m r^2 \dot{\theta})} \hat{e}_\theta$$

$$+ \frac{d}{dt}(\gamma m \dot{z}) \hat{e}_z$$

(NOTE: ON THIS PAGE $p_\theta^* \equiv \theta$-component of MECHANICAL MOMENTUM

NOT TO BE CONFUSED WITH $P_\theta = \gamma m r^2 \dot{\theta} + q r A_\theta \equiv \theta$-component

OF <u>CANONICAL</u> <u>ANGULAR</u> MOMENTUM)

CONSERVATION OF CANONICAL ANGULAR MOMENTUM

DEFINE $p_\theta = \gamma m r^2 \dot{\theta} + q r A_\theta$

$$\frac{d}{dt} p_\theta = 0$$ (CONSERVATION OF CANONICAL ANGULAR MOMENTUM)

NOTE THAT THE FLUX ENCLOSED BY A CIRCLE OF RADIUS r

$$\psi = \int \underline{B} \cdot d\underline{A} = \int (\nabla \times \underline{A}) \cdot d\underline{A} = \oint \underline{A} \cdot d\underline{l} = 2\pi r A_\theta$$

$$p_\theta = \gamma m r^2 \dot{\theta} + \frac{q}{2\pi} \psi$$ IS CONSERVED ALONG AN ORBIT IN AXISYMMETRIC GEOMETRIES

"EXTERNAL" (REISER SECTION 3.3)

ELECTRIC & MAGNETIC FIELDS WITH RADIAL SYMMETRY

J. BALMAIN

(B)

CONSIDER THE EXTERNAL FIELD: (TIME STEADY VACUUM SOLUTIONS)

$$\nabla \times \underline{B} = 0 \quad \& \quad \nabla \times \underline{E} = 0$$

$$\nabla \cdot \underline{B} = 0 \quad \& \quad \nabla \cdot \underline{E} = 0 \quad \Rightarrow \quad \underline{E} \ \& \ \underline{B} = -\underline{\nabla}\Phi$$

Let $\Phi = \sum_{\nu=0}^{\infty} f_{2\nu}(z) r^{2\nu}$ $\quad \nabla^2 \phi = 0 \Rightarrow \Phi = \sum_{\nu=0}^{\infty} \frac{(-1)^{\nu}}{\nu!} \frac{\partial^{2\nu} f(0,z)}{\partial z^{2\nu}} \left(\frac{r}{2}\right)^{2\nu}$

$$\Rightarrow \Phi = \Phi(0,z) - \frac{1}{4}\frac{\partial^2 \Phi(0,z)}{\partial z^2} r^2 + \frac{1}{64}\frac{\partial^4 \Phi(0,z)}{\partial z^4} r^4$$

Let $B_z(0,z) = B(z)$ & let $\Phi(0,z) = V(z)$

$$B_z(r,z) = B(z) - \frac{r^2}{4}\frac{\partial^2 B}{\partial z^2} + \frac{r^4}{64}\frac{\partial^4 B}{\partial z^4} + \cdots$$

$$B_r(r,z) = -\frac{r}{2}\frac{\partial B}{\partial z} + \frac{r^3}{16}\frac{\partial^3 B}{\partial z^3} + \cdots$$

$$\Phi(r,z) = V(z) - \frac{1}{4}V'' r^2 + \frac{r^4}{64}\frac{\partial^4 V}{\partial z^4}$$

$$\Rightarrow \quad E_r = \frac{1}{2}V'' r - \frac{r^3}{16}\frac{\partial^4 V}{\partial z^4}$$

Also, $\psi \cong \pi r^2 B(z)$

$$\nabla^2 \phi = \frac{1}{r}\frac{\partial}{\partial r}\left(r\frac{\partial \phi}{\partial r}\right) + \frac{\partial^2 \phi}{\partial z^2} = 0$$

$$f(r,z) = \sum_{\nu=0}^{\infty} f_{2\nu}(z)\, r^{2\nu} = f_0 + f_2 r^2 + f_4 r^4$$

$$\sum_{\nu=1}^{\infty} \left[2\nu + 2\nu(\nu-1)\right] f_{2\nu}\, r^{2\nu-2} + \sum_{\nu=0}^{\infty} f_{2\nu}''\, r^{2\nu} = 0$$

PARAXIAL RAY EQUATION

$$(\mathrm{I}) \Rightarrow \underbrace{\frac{d}{dt}(\gamma m \dot{r})}_{\text{INERTIAL}} - \underbrace{\gamma m r \dot{\theta}^2}_{\text{CENTRIFUGAL}} = q\Big(\underbrace{\frac{V''}{2} r}_{E_r \text{ external}} + \underbrace{r\dot{\theta} B}_{v_\theta B_z \text{ external}}\Big) + q\Big(\underbrace{E_r^{self} - v_z B_\theta^{self}}_{\text{SELF FIELDS}}\Big)$$

Now use s as independent variable $v_z dt = ds$

$$v_z \frac{d}{ds}\left(\gamma m v_z r'\right) - \gamma m v_z^2 r \theta'^2 = q\left(\frac{V''}{2} r + r v_z \theta' B\right) + q\left(E_r^{self} - v_z B_\theta^{self}\right)$$

EXPANDING 1st term and $v_z \cong v$; AND DIVIDING BY $\gamma m v^2$:

$$r'' - r\theta'^2 + \frac{\gamma'}{\beta^2 \gamma} r' = \frac{q}{\gamma m \beta^2 c^2}\left(\frac{V''}{2} r + r\beta c \theta' B + E_r^{self} - v_z B_\theta^{self}\right) \qquad (P1)$$

Using CANONICAL MOMENTUM, eliminate θ' via

$$\theta' = \frac{p_\theta - \frac{q\psi}{2\pi}}{\gamma m r^2 \beta c} = \frac{p_\theta}{\gamma m r^2 \beta c} - \frac{qB}{2\gamma \beta m c} \equiv \frac{p_\theta}{\gamma m r^2 \beta c} - \frac{\omega_c}{2\gamma\beta c}$$

where we define $\omega_c \equiv \frac{qB}{m}$.

ADDING THE TWO θ' TERMS IN THE EQUATION (P1)

$$-r\theta'^2 - \frac{r\omega_c \theta'}{\gamma\beta c} = \frac{-p_\theta^2}{\gamma^2 m^2 r^3 \beta^2 c^2} + \frac{p_\theta \omega_c}{\gamma^2 m \beta^2 c^2 r} - \frac{r\omega_c^2}{4\gamma^2\beta^2 c^2} - \frac{p_\theta \omega_c}{\gamma^2 m \beta^2 c^2 r} + \frac{r\omega_c^2}{2\gamma^2\beta^2 c^2}$$

$$= \frac{-p_\theta^2}{\gamma^2 m^2 r^3 \beta^2 c^2} + \frac{r\omega_c^2}{2\gamma^2\beta^2 c^2}$$

So equation (P1) becomes:

$$r'' + \frac{\gamma'}{\beta^2\gamma} r' = \frac{q}{\gamma m \beta^2 c^2}\left(\frac{V''}{2} r\right) + \frac{r\omega_c^2}{2\gamma^2\beta^2c^2} + \frac{p_\theta^2}{\gamma^2 m^2 r^3 \beta^2 c^2} + \frac{q}{\gamma m \beta^2 c^2}\left[E_r^{self} - v_z B_\theta^{self}\right] \quad (P2)$$

Now $\gamma' mc^2 = q\frac{\underline{E}\cdot\underline{v}}{v_z}$ so $\gamma'' = \left(V'' + \frac{\partial^2\phi^{self}}{\partial z^2}\right)\frac{q}{mc^2}$

CALCULATING $\frac{q}{\gamma m\beta^2c^2}\left[\frac{V''}{2}r + E_r^{self} - v_z B_\theta^{self}\right]$:

$$\nabla^2\phi^{self} = -\frac{\rho}{\varepsilon_0} \Rightarrow \frac{1}{r}\frac{\partial}{\partial r}\left(r\frac{\partial\phi}{\partial r}\right) = -\frac{\rho}{\varepsilon_0} - \frac{\partial^2\phi^{self}}{\partial z^2}$$

$$\Rightarrow \frac{\partial}{\partial r}\left(r\frac{\partial\phi}{\partial r}\right) = -\frac{r\rho(r)}{\varepsilon_0} - \frac{r\partial^2\phi^{self}}{\partial z^2}$$

$$r\frac{\partial\phi}{\partial r} = -\frac{1}{2\pi\varepsilon_0}\int_0^r 2\pi r\,\rho(r)\,dr - \frac{r^2}{2}\frac{\partial^2\phi}{\partial z^2}$$

$$= -\frac{1}{2\pi\varepsilon_0}\lambda(r) - \frac{r^2}{2}\frac{\partial^2\phi^{self}}{\partial z^2}$$

$$\Rightarrow E_r^{self} = \frac{\lambda(r)}{2\pi\varepsilon_0 r} + \frac{r}{2}\frac{\partial^2\phi^{self}}{\partial z^2}$$

$$\nabla\times\underline{B} = \mu_0\underline{J} \Rightarrow 2\pi r B_\theta = \mu_0\int_0^r 2\pi r\, J_z(r)\,dr = \mu_0 v_z\lambda(r)$$

$$B_\theta^{self} = \frac{\mu_0 v_z\lambda(r)}{2\pi r} = \frac{v_z}{c^2}\frac{\lambda(r)}{2\pi\varepsilon_0 r}$$

$$\left[\frac{V''}{2}r + E_r^{self} - v_z B_\theta^{self}\right] = \left[\frac{r}{2}\underbrace{\left(V'' + \frac{\partial^2\phi^{self}}{\partial z^2}\right)}_{-\frac{mc^2}{q}\gamma''} + \underbrace{\left(1 - \frac{v_z^2}{c^2}\right)}_{1/\gamma^2}\frac{\lambda(r)}{2\pi\varepsilon_0 r}\right]$$

So equation (P2) becomes: "THE PARAXIAL RAY EQUATION:"

$$\underbrace{r'' + \frac{\gamma'}{\beta^2\gamma} r'}_{\text{INERTIAL}} + \underbrace{\frac{\gamma''}{2\beta^2\gamma} r}_{\substack{E_r \\ \text{(CONVERGENCE OF FIELD LINES)}}} + \underbrace{\left(\frac{\omega_c}{2\gamma\beta c}\right)^2 r}_{\substack{V_\theta B_z \\ \text{- CENTRIFUGAL}}} - \underbrace{\left(\frac{p_\theta}{\gamma\beta m c}\right)^2 \frac{1}{r^3}}_{\text{CENTRIFUGAL}} - \underbrace{\frac{q}{\gamma^3 m v_z^2} \frac{\lambda(r)}{2\pi\epsilon_0 r}}_{\text{SELF FIELD}} = 0$$

MOMENT EQUATIONS

Vlasov eqtn: $\frac{\partial f}{\partial s} + x'\frac{\partial f}{\partial x} + x''\frac{\partial f}{\partial x'} + y'\frac{\partial f}{\partial y} + y''\frac{\partial f}{\partial y'} = 0$

Let $g = g(x, x', y, y')$; $N = \iiiint f \, dx\, dx'\, dy\, dy'$

MULTIPLY VLASOV equation by g & $\frac{1}{N}\iiiint dx\, dx'\, dy\, dy'$

$$\int dx\, dx'\, dy\, dy \left[g\frac{\partial f}{\partial s} + g x'\frac{\partial f}{\partial x} + g x''\frac{\partial f}{\partial x'} + g y'\frac{\partial f}{\partial y} + g y''\frac{\partial f}{\partial y'} \right] = 0$$

$$\Rightarrow \frac{d}{ds}\langle g \rangle + \overbrace{\underbrace{\frac{1}{N}\iiint g f \Big|_{x=-\infty}^{\infty}}_{\to 0} - \underbrace{\frac{1}{N}\iiiint \frac{\partial g}{\partial x} f x'}_{= \langle x' \frac{\partial g}{\partial x} \rangle}}^{\text{INTEGRATE BY PARTS}} + \cdots = 0$$

$$\Rightarrow \frac{d}{ds}\langle g \rangle = \langle x'\frac{\partial g}{\partial x} \rangle + \langle x''\frac{\partial g}{\partial x'} \rangle + \langle y'\frac{\partial g}{\partial y} \rangle + \langle y''\frac{\partial g}{\partial y'} \rangle$$

But $\frac{dg}{ds} = \frac{\partial g}{\partial x}x' + \frac{\partial g}{\partial x'}x'' + \frac{\partial g}{\partial y}y' + \frac{\partial g}{\partial y'}y''$

$$\Rightarrow \frac{d}{ds}\langle g \rangle = \langle g' \rangle$$

So $\frac{d}{ds}\langle x^2 \rangle = 2\langle x x' \rangle$

$\frac{d}{ds}\langle x'^2 \rangle = 2\langle x' x'' \rangle$ etc...

$\frac{d}{ds}\langle x x' \rangle = \langle x x'' \rangle + \langle x'^2 \rangle$

ENVELOPE EQUATION FOR AXISYMMETRIC BEAMS

LET $r_b^2 = 2\langle r^2\rangle = 2(\langle x^2\rangle + \langle y^2\rangle) = 4\langle x^2\rangle$ for an axisymmetric beam

$$2 r_b r_b' = 4\langle r r'\rangle \quad \Rightarrow \quad r_b' = \frac{2\langle r r'\rangle}{r_b}$$

$$r_b'' = \frac{2\langle r r''\rangle + 2\langle r'^2\rangle}{r_b} - \frac{2\langle r r'\rangle}{r_b^2}\left(\frac{2\langle r r'\rangle}{r_b}\right)$$

$$= 2\frac{\langle r r''\rangle}{r_b} + \frac{4\langle r^2\rangle\langle r'^2\rangle - 4\langle r r'\rangle^2}{r_b^3}$$

WHAT IS $\langle r r''\rangle$?

RECALL EQUATION P1 (ON PATH TO PARAXIAL RAY EQUATION):

$$r'' - r\theta'^2 + \frac{\gamma'}{\beta^2\gamma} r' = \frac{q}{\gamma m \beta^2 c^2}\left(\frac{V''}{2} r + r\beta c\theta' B + E_r^{self} - v_z B_z^{self}\right)$$

P1 may be rewritten:

$$r'' - r\theta'^2 + \frac{\gamma'}{\beta^2\gamma} r' = \frac{q}{\gamma m \beta^2 c^2}\left[\frac{-mc^2}{q}\gamma''\frac{r}{2} + \frac{\lambda(r)}{\gamma^2 2\pi\epsilon_0 r} + r\beta c\theta' B\right]$$

$\Rightarrow$

$$\boxed{r'' + \frac{\gamma'}{\beta^2\gamma} r' + \frac{\gamma''}{2\beta^2\gamma} r - \frac{q}{\gamma^3 m v_z^2}\frac{\lambda(r)}{2\pi\epsilon_0 r} \underbrace{\frac{-\omega_c}{\gamma\beta c}\theta' r - r\theta'^2}_{} = 0}$$

What is $\langle r r'' \rangle$?

$$\langle r r'' \rangle + \frac{-\omega_c}{\gamma\beta c}\langle \theta' r^2 \rangle - \langle r^2\theta'^2 \rangle + \cdots = 0$$

$$\langle p_\theta \rangle^2 = \gamma^2 m^2 \beta^2 c^2 \langle r^2\theta' \rangle^2 + \frac{\omega_c^2}{4} m^2 \langle r^2 \rangle^2 + \omega_c \gamma m^2 \beta c \langle r^2\theta' \rangle \langle r$$

$$\Rightarrow \frac{-\omega_c}{\gamma\beta c}\langle \theta' r^2 \rangle = \frac{-\omega_c}{\gamma\beta c}\left[\frac{\langle p_\theta \rangle^2}{\omega_c \gamma m^2 \beta c \langle r^2 \rangle} - \frac{\omega_c \langle r^2 \rangle}{4\gamma\beta c} - \frac{\gamma\beta c}{\omega_c}\frac{\langle r^2\theta' \rangle^2}{\langle r^2 \rangle}\right]$$

$$\Rightarrow \langle r r'' \rangle = \frac{\langle p_\theta \rangle^2}{\gamma^2 m^2 \beta^2 c^2 \langle r^2 \rangle} - \frac{\omega_c^2 \langle r^2 \rangle}{4\gamma^2 \beta^2 c^2} - \frac{\langle r^2\theta' \rangle^2}{\langle r^2 \rangle} + \langle r^2\theta'^2 \rangle + \cdots = 0$$

$\Rightarrow$

$$\langle r r'' \rangle = \frac{\gamma'}{\beta^2 \gamma} \langle r r' \rangle + \frac{\gamma''}{2\beta^2 \gamma} \langle r^2 \rangle - \frac{q}{\gamma^3 m v_z^2} \frac{\langle \lambda(r) \rangle}{2\pi \epsilon_0} +$$

$$\frac{\langle p_\theta \rangle^2}{(\gamma m \beta c)^2 \langle r^2 \rangle} - \frac{\omega_c^2 \langle r^2 \rangle}{4(\gamma^2 \beta c)^2} - \frac{\langle r^2 \theta' \rangle^2}{\langle r^2 \rangle} + \langle r^2 \theta'^2 \rangle$$

$$r_b'' = \frac{2\langle r r'' \rangle}{r_b} + \frac{4\langle r^2 \rangle \langle r'^2 \rangle - 4\langle r r' \rangle}{r_b^3}$$

$$= \frac{\gamma'}{\beta^2 \gamma} \frac{2\langle r r' \rangle}{r_b} + \frac{\gamma''}{2\beta^2 \gamma} \frac{2\langle r^2 \rangle}{r_b} - \frac{2q}{\gamma^3 m v_z^2} \frac{\langle \lambda(r) \rangle}{2\pi \epsilon_0} \frac{1}{r_b}$$

$$+ \frac{\langle p_\theta \rangle^2}{(\gamma m \beta c)^2} \frac{2}{\langle r^2 \rangle r_b} - \frac{\omega_c^2}{4(\gamma \beta c)^2} \frac{2\langle r^2 \rangle}{r_b} - \frac{2\langle r^2 \theta' \rangle^2}{r_b \langle r^2 \rangle}$$

$$+ \frac{2\langle r^2 \theta'^2 \rangle}{r_b} + \frac{4\langle r^2 \rangle \langle r'^2 \rangle - 4\langle r r' \rangle^2}{r_b^3}$$

USING $r_b^2 \equiv 2\langle r^2 \rangle$ & $r_b' = \dfrac{2\langle r r' \rangle}{r_b}$

ENVELOPE EQUATION

$$\Rightarrow \boxed{r_b'' + \frac{\gamma'}{\beta^2 \gamma} r_b' + \frac{\gamma''}{2\beta^2 \gamma} r_b + \left(\frac{\omega_c}{2\gamma \beta c}\right)^2 r_b + \frac{-4\langle p_\theta \rangle^2}{(\gamma m \beta c)^2 r_b^3} - \frac{\varepsilon_r^2}{r_b^3} - \frac{Q}{r_b} = 0}$$

WHERE $\varepsilon_r^2 = 4(\langle r^2 \rangle \langle r'^2 \rangle - \langle r r' \rangle^2 + \langle r^2 \rangle \langle r^2 \theta'^2 \rangle - \langle r^2 \theta' \rangle^2)$

ENVELOPE EQUATION -- CONTINUED

$$\boxed{r_b'' + \frac{\gamma'}{\beta^2\gamma} r_b' + \frac{\gamma''}{2\beta^2\gamma} r_b + \left(\frac{\omega_c}{2\gamma\beta c}\right)^2 r_b - \frac{4\langle p_\theta\rangle^2}{(\gamma m\beta c)^2 r_b^3} - \frac{\varepsilon_r^2}{r_b^3} - \frac{Q}{r_b} = 0}$$

COMPARE WITH THE SINGLE PARTICLE PARAXIAL RAY EQUATION:

$$\underbrace{r'' + \frac{\gamma'}{\beta^2\gamma} r'}_{\text{INERTIAL}} + \underbrace{\frac{\gamma''}{2\beta^2\gamma} r}_{E_r} + \underbrace{\left(\frac{\omega_c}{2\gamma\beta c}\right)^2 r}_{v_\theta B_z - \text{CENTRIFUGAL}} - \underbrace{\left(\frac{p_\theta}{\gamma m\beta c}\right)^2 \frac{1}{r^3}}_{\text{CENTRIFUGAL}} - \underbrace{\frac{q}{\gamma^3 m v_z^2} \frac{\lambda(r)}{2\pi\varepsilon_0 r}}_{E_r - v_z B_\theta \text{ self field}} = 0$$

$$\varepsilon_r^2 = 4\left(\langle r^2\rangle\langle r'^2\rangle - \langle rr'\rangle^2 + \langle r^2\rangle\langle r^2\theta'^2\rangle - \langle r^2\theta'\rangle^2\right)$$

NOTE THAT FOR AXISYMMETRIC BEAMS ($\rho = \rho(r)$ ONLY)

$$\langle r^2\rangle = \langle x^2\rangle + \langle y^2\rangle = 2\langle x^2\rangle$$

$$\Rightarrow 2\langle rr'\rangle = 4\langle xx'\rangle$$

$$\& \quad \langle x'^2\rangle + \langle y'^2\rangle = 2\langle x'^2\rangle = \langle r'^2\rangle + \langle r^2\theta'^2\rangle$$

DEFINE $\varepsilon_x^2 = 16\left(\langle x^2\rangle\langle x'^2\rangle - \langle xx'\rangle^2\right)$

$$\Rightarrow \boxed{\varepsilon_r^2 = \varepsilon_x^2 - 4\langle r^2\theta'\rangle^2}$$

EXAMPLES OF
SYSTEMS WITH AXIAL SYMMETRY

- PERIODIC SOLENOIDS

- EINZEL LENSES

- CONTINUOUS FOCUSING

EXAMPLES OF
SYSTEMS WITHOUT AXIAL SYMMETRY

- ELECTRIC OR MAGNETIC QUADRUPOLE

⟹ USE CARTESIAN COORDINATES WITH
ELLIPTICAL SPACE CHARGE SYMMETRY

TOPS
35500

(11)



Figure 3.3. Schematic of conductor configuration with applied voltages producing an alternating-gradient quadrupole electric field with axial periodicity length S.



Figure 3.7. Alternating step-function model of a periodic quadrupole lattice with filling factor η for the lens elements. The figure shows a plot of the quadrupole coupling coefficient $\kappa_q(s)$ versus s for one full period (S) of the lattice. Such a configuration is often called a FODO transport lattice (acronym for focusing-off-defocusing-off).

FIGURES FROM DAVIDSON & QIN 2003

(12)

figure from
Davidson & Qin, 2003.



Figure 3.1. Schematic of magnet sets producing an alternating-gradient quadrupole field with axial periodicity length S.

J. BARNARD
13

FROM REISER, p. 112

$E_x = -E'x$
$E_y = E'y$



$F_x = -qE'x$
$F_y = qE'y$

ELECTROSTATIC QUADS

Figure 3.15. Electrodes and force lines in an electrostatic quadrupole.

$B_x = B'y$
$B_y = B'x$

$F_x = -qV_z B'x$
$F_y = qV_x B'y$



MAGNETIC QUADS

QUADRUPOLE FOCUSING

Now, relax radial symmetry:

For $\nabla \cdot B = 0$ & $\nabla \times B = 0$

EXPAND FIELD IN CYLINDRICAL "MULTIPOLES":

$$E_r, B_r = \sum_{n=1}^{\infty} f_n r^{n-1} \cos(n\theta)$$



$E_x = E_r \cos\theta - E_\theta \sin\theta$
$E_y = E_r \sin\theta + E_\theta \cos\theta$

$$E_\theta, B_\theta = \sum_{n=1}^{\infty} f_n r^{n-1} \sin(n\theta)$$

$n=1 \Rightarrow$ dipole $\begin{cases} E_r = f_1 \cos\theta \\ E_\theta = -f_1 \sin\theta \end{cases} \Rightarrow \begin{matrix} E_x = f_1 \\ E_y = 0 \end{matrix}$

$n=2 \Rightarrow$ quadrupole $\begin{cases} E_r = f_2 r \cos 2\theta \\ E_\theta = -f_2 r \sin 2\theta \end{cases} \Rightarrow \begin{matrix} E_x = f_2 x \\ E_y = -f_2 y \end{matrix}$

NOTE: ABOVE EXPANSION IS VALID WHEN <u>E or B ≠ function(z)</u>. FOR MAGNETS OF FINITE AXIAL EXTENT, FOR EACH FUNDAMENTAL n-pole, A SET OF HIGHER ORDER MULTIPOLES WITH SAME AZIMUTHAL SYMMETRY ARE REQUIRED TO SATISFY $\nabla^2\phi = 0$.

FOR EXAMPLE FOR A FUNDAMENTAL QUADRUPOLE THE FIELD MAY BE EXPANDED:

$$E_r = \sum_{\nu=0}^{\infty} f_{2,\nu}(z)\,[1+\nu]\, r^{1+2\nu} \cos[2\theta]$$

$$E_\theta = \sum_{\nu=0}^{\infty} -f_{2,\nu}(z)\, r^{1+2\nu} \sin[2\theta]$$

$$E_z = \sum_{\nu=0}^{\infty} \frac{1}{2}\frac{df_{2,\nu}}{dz}\, r^{2+2\nu} \cos 2\theta$$

with $f_{2,\nu+1}(z) = \dfrac{-1}{4(\nu+1)(\nu+3)} \dfrac{d^2 f_{2,\nu}(z)}{dz^2}$

SEE LUND, S. M. (1996) FOR EXAMPLE. HIF note 96-1 LLNL.

Heavy ion accelerators use alternating gradient quadrupoles to focus (confine) the beams (non-neutral plasmas)

Space-charge forces and thermal forces act to expand beam



Magnetic quad



Electric quad

Quadrupoles (magnetic or electric):

- alternately provide inward then outward impulse
- focus in one plane and defocus in other
- act as linear lenses. (Force proportional to distance from axis).

Horizontal (x) plane:



Vertical (y) plane:



Average displacement is larger in focusing lenses so the net effect is focusing.

J. Barnard

16

Space charge reduces betatron phase advance



J. BARNARD

17