John Barnard
Steven Lund
USPAS
June 2006

Summary of JB lectures

START WITH MICROSCOPIC PHASE SPACE DENSITY

$$\boxed{N(\underline{x},\underline{v},t) = \sum_{i=1}^{N} \delta(\underline{x}-\underline{X}_i(t))\,\delta(\underline{v}-\underline{V}_i(t))}$$ Klimontovich Density

$\frac{\partial N}{\partial t}$ + LORENTZ EQUATIONS $\Rightarrow$ KLIMONTIVICH EQUATION:

$$\boxed{\frac{\partial N}{\partial t} + v\cdot\nabla_x N(\underline{x},\underline{v},t) - \frac{q}{m}(E^m + v\times B^m)\cdot\nabla_v N(\underline{x},\underline{v},t) = 0}$$

$$\boxed{\text{or } \frac{dN(\underline{x},\underline{v},t)}{dt} = 0}$$

Letting $N = f + \delta f$ $\quad f = \langle N\rangle$ $\quad f = \frac{\int N\, d^3x\, d^3v}{\Delta x^3\,\Delta v^3}$

$E^m = E + \delta E$ $\quad E = \langle E_m\rangle$

$B^m = B + \delta B$ $\quad B = \langle B_m\rangle$ $\quad n^{-1/3} \ll \Delta x \ll \lambda_D$

PERFORMING LOCAL AVERAGES TO OBTAIN SMOOTH & "SLIKEY" QUANTITIES:

$$\boxed{\frac{\partial f}{\partial t} + \underline{v}\cdot\frac{\partial f}{\partial \underline{x}} + \frac{d\underline{v}}{dt}\cdot\frac{\partial f}{\partial \underline{v}} = \frac{\partial f}{\partial t_c}} \sim \frac{f}{\tau_c}$$

We estimated $\frac{|\partial f/\partial t_c|}{\left|\frac{q\underline{E}}{m}\cdot\partial f/\partial \underline{v}\right|} \sim \frac{1}{16\,\lambda_D^3\, n_0} \ll 1$

$\lambda_D = v_{TH}/\omega_p \qquad v_{TH} \equiv \sqrt{\frac{kT}{m}} \qquad \omega_p \equiv \sqrt{\frac{q^2 n}{\varepsilon_0 m}}$

$$\frac{\partial f}{\partial t} + \underline{v}\cdot\frac{\partial f}{\partial \underline{x}} + \frac{d\underline{p}}{dt}\cdot\frac{\partial f}{\partial \underline{p}} = 0 \qquad \dot{p} = -\frac{\partial H}{\partial \underline{x}}\ ;\ \dot{x} = \frac{\partial H}{\partial \underline{p}}$$

$$\boxed{\frac{df}{dt} = 0}$$ LIOUVILLE'S EQUATION (INCOMPRESSIBILITY OF PHASE VOLUME)

DEFINE NORMALIZED EMITTANCES PROPORTIONAL TO $\Delta p_z \Delta z \propto \Delta E\,\Delta t$

$\Delta p_x \Delta x$

$\Delta p_y \Delta y$

SO THAT

$$\varepsilon_{Nx}^2 = \gamma^2\beta^2\left(\langle x^2\rangle\langle x'^2\rangle - \langle x x'\rangle^2\right)$$

$\Rightarrow$ CONSTANT IF FORCES ARE LINEAR IN X & FILAMENTATION IS ABSENT. (LINEAR WITHOUT COUPLING TO Z, OR Y).

WE DERIVED TWO SETS OF PARTICLE EQUATION OF MOTION:

PARAXIAL EQUATION (FOR AXISYMMETRIC SYSTEMS) $\left(\frac{\partial}{\partial\theta} = 0\right)$

STARTING WITH THE LORENTZ FORCE EQUATION $\frac{d\underline{p}}{dt} = q(\underline{E} + \underline{v} \times \underline{B})$ IN CYL. COORD.

$$\underbrace{\frac{d}{dt}(\gamma m \dot{r})}_{\text{INERTIAL}} - \underbrace{\gamma m r \dot{\theta}^2}_{\text{CENTRIFUGAL}} = q\left(\underbrace{\frac{V''}{2} r}_{E_r \text{ external (DIVERGENCE OF } E = 0)} + \underbrace{r\dot{\theta} B}_{v_\theta B_z}\right) + q\left(\underbrace{E_r^{self}}_{\text{SELF-FIELD}} + \underbrace{v_z B_\theta^{self}}_{\text{SELF-FIELD}}\right)$$

$$\boxed{\dot{r} \equiv \frac{dr}{dt}; \quad r' = \frac{dr}{ds} = \frac{\dot{r}}{\beta c}}$$

θ-component:

$$p_\theta = \gamma m r^2 \dot{\theta} + \frac{qB(z) r^2}{2} = \text{constant}$$

$$= \gamma m r^2 \beta c \theta' + \frac{qBr^2}{2} = \text{constant}$$

$$\underbrace{r''}_{\text{INERTIAL}} + \underbrace{\frac{\gamma'}{\beta^2\gamma} r'}_{\text{ACCELERATION (INERTIA)}} + \underbrace{\frac{\gamma''}{2\beta^2\gamma} r}_{E_r \text{ (CONVERGENCE OF FIELD LINES)}} + \underbrace{\left(\frac{\omega_c}{2\gamma\beta c}\right)^2 r}_{v_\theta B_z \text{ CENTRIFUGAL}} - \underbrace{\left(\frac{p_\theta}{\gamma\beta m c}\right)^2 \frac{1}{r}}_{\text{CENTRIFUGAL}} - \underbrace{\frac{q}{\gamma^3 m v_z^2} \frac{\lambda(r)}{2\pi\epsilon_0 r}}_{\text{SELF FIELD}} = 0$$

STATISTICAL AVERAGE OF THIS EQUATION

$$r_b'' + \frac{\gamma'}{\beta^2\gamma} r_b' + \frac{\gamma''}{2\beta^2\gamma} r_b + \left(\frac{\omega_c}{2\gamma\beta c}\right)^2 r_b - \frac{4\langle p_\theta\rangle^2}{(\gamma m\beta c)^2 r_b^3} - \frac{\varepsilon_r^2}{r_b^3} - \frac{Q}{r_b} = 0$$

$$\varepsilon_r^2 \equiv 4\left(\langle r^2\rangle\langle r'^2\rangle - \langle r r'\rangle^2 + \langle r^2\rangle\langle r^2\theta'^2\rangle - \langle r^2\theta'\rangle^2\right); \quad Q = \frac{q\lambda}{2\pi\epsilon_0 \gamma^3 \beta^2 m c^2}$$

$$= \varepsilon_x^2 - 4\langle r^2\theta'\rangle^2 \quad (\text{if } \rho = \rho(r) \text{ only})$$

EQUATION OF MOTION

AGAIN STARTING WITH $\frac{d\underline{p}}{dt} = q(\underline{E} + \underline{v} \times \underline{B})$

RETURN TO X, Y COORDINATES

$$x'' + \frac{1}{\gamma v_z}\frac{d}{ds}(\gamma v_z)x' = \frac{-q}{\gamma^3 m v_z^2}\frac{\partial \varphi}{\partial x} \pm \begin{cases} \frac{qB'}{\gamma m v_z} x & \text{for magnetic quads} \\ \frac{qE'}{\gamma m v_z^2} x & \text{for electric quads} \end{cases}$$

Let $\frac{\gamma m v_z}{q} = \frac{p}{q} \equiv [B\rho] \equiv$ RIGIDITY

$$y'' + \frac{1}{\gamma v_z}\frac{d}{ds}(\gamma v_z)y' = \frac{-q}{\gamma^3 m v_z^2}\frac{\partial \varphi}{\partial y} \mp \begin{cases} \frac{B'}{[B\rho]} y & \text{magnetic} \\ \frac{qE'}{\gamma m v_z^2} y & \text{electric} \end{cases}$$

ENVELOPE EQUATION

$$r_x^2 = 4\langle x^2\rangle; \quad r_y^2 = 4\langle y^2\rangle$$

$$r_x' = \frac{4\langle x x'\rangle}{r_x}$$

$$r_x'' = \frac{4\langle x x''\rangle}{r_x} + \frac{\varepsilon_x^2}{r_x^3}; \qquad \varepsilon_x^2 = 16(\langle x^2\rangle\langle x'^2\rangle - \langle x x'\rangle^2)$$

$$r_y'' = \frac{4\langle y y''\rangle}{r_y} + \frac{\varepsilon_y^2}{r_y^3} \qquad \varepsilon_y^2 = 16(\langle y^2\rangle\langle y'^2\rangle - \langle y y'\rangle^2)$$

for magnetic focusing:

$$r_x'' + \frac{1}{\gamma v_z}\frac{d}{ds}(\gamma v_z) r_x' + \frac{4q}{\gamma^3 m v_z^2}\frac{\langle x \frac{\partial \varphi}{\partial x}\rangle}{r_x} \pm \frac{B'}{[B\rho]} r_x - \frac{\varepsilon_x^2}{r_x^3} = 0$$

$$r_y'' + \frac{1}{\gamma v_z}\frac{d}{ds}(\gamma v_z) r_y' + \frac{4q}{\gamma^3 m v_z^2}\frac{\langle y \frac{\partial \varphi}{\partial y}\rangle}{r_y} \pm \frac{B'}{[B\rho]} r_x - \frac{\varepsilon_y^2}{r_y^3} = 0$$

(for electric focusing $\frac{B'}{[B\rho]} \to \frac{qE'}{\gamma m v_z^2}$)

National® Brand 42-182 100 SHEETS Made in U.S.A.

SPACE CHARGE TERM WITH ELLIPTICAL SYMMETRY

NOW DEFOCUSING IN ONE DIRECTION AND FOCUSING IN THE OTHER $\Rightarrow$ RADIAL SYMMETRY SHOULD BE REPLACED BY ELLIPTICAL SYMMETRY: $\rho = \rho\left(\frac{x^2}{r_x^2} + \frac{y^2}{r_y^2}\right)$

CAN BE SHOWN THAT $\left\langle x \frac{\partial \phi}{\partial x} \right\rangle = \frac{-\lambda}{4\pi\epsilon_0} \frac{r_x}{r_x + r_y}$

$$\left\langle y \frac{\partial \phi}{\partial y} \right\rangle = \frac{-\lambda}{4\pi\epsilon_0} \frac{r_y}{r_x + r_y}$$

USE $\phi(x,y) = \frac{-r_x r_y}{4\epsilon_0} \int_0^\infty \frac{\hat{\rho}(\chi)\, ds}{\sqrt{r_x^2+s}\sqrt{r_y^2+s}}$ to prove, where $\hat{\rho}(\chi) = \frac{d\rho}{d\chi}$

$\rho(x,y) = \hat{\rho}(\chi)\big|_{s=0}$

$\chi = \frac{x^2}{r_x^2+s} + \frac{y^2}{r_y^2+s}$

DEFINING $Q = \frac{2\lambda q}{4\pi\epsilon_0 \gamma^3 m v_z^2}$

$$\Rightarrow \boxed{\begin{aligned} & r_x'' + \frac{1}{\gamma v_z}\frac{d}{ds}(\gamma v_z) r_x' - \frac{2Q}{r_x + r_y} \mp \frac{B'}{[B\rho]} r_x - \frac{\varepsilon_x^2}{r_x^3} = 0 \\ & r_y'' + \frac{1}{\gamma v_z}\frac{d}{ds}(\gamma v_z) r_y' - \frac{2Q}{r_x + r_y} \pm \frac{B'}{[B\rho]} r_y - \frac{\varepsilon_y^2}{r_y^3} \end{aligned}}$$

(for Electric Focusing $\frac{B'}{[B\rho]} \to \frac{qE'}{\gamma m v_z^2}$)

(ANALOGUE TO CIRCULAR BEAM:

$\left\langle r \frac{\partial \phi}{\partial r} \right\rangle = \frac{-\lambda}{4\pi\epsilon_0}$ PROVED IN HOMOWORK)

ENVELOPE EQUATIONS DERIVED

RMS (ENVELOPES DEFINED IN TERMS OF RMS QUANTITIES; EMITTANCE NOT GUARANTEED TO BE CONSERVED).

1. PARAXIAL r_b ; $\rho(r)$
2. ELLIPTICAL r_x, r_y ; $\rho\left(\frac{x^2}{r_x^2}+\frac{y^2}{r_y^2}\right)$
3. LONGITUDINAL r_z FOR $E_z = \frac{-q}{4\pi\varepsilon_0}\frac{\partial\lambda}{\partial z} \propto z$; $[\lambda \propto (1-4z^2/r_z^2)]$
   $V \propto (z/r_z)$
4. ELLIPSOIDAL of BUNCHES $r_\perp, r_z$ (ALSO r_x, r_y, r_z c.f. Wangler, section 9.9).
   (ρ CONSTANT)
5. ELLIPTICAL WITH IMAGES r_x, r_y $\rho\left(\frac{x^2}{r_x^2}+\frac{y^2}{r_y^2}\right)$
6. BARRIER FRAME (PERIODIC SOLENOIDS OR CONTINUOUS FOCUSING) $\tilde{r}_x, \tilde{r}_y$ $\rho(r)$

KINETIC ENVELOPE EQUATIONS (CONSTRAINT EQUATION OF A PARTICULAR DISTRIBUTION FUNCTION; EMITTANCE CONSERVED)

1. KV FOR ELLIPTICAL UNIFORM BEAMS $f(x,x',y,y')$
   [IDENTICAL TO #2 ABOVE]
2. NEUFFER DISTRIBUTION FOR 1D $f(z,z')$
   [IDENTICAL TO #3 ABOVE]

MOMENT EQUATIONS

1. TRANSVERSE WITH CHROMATIC EFFECTS
   $\langle x^2\rangle, \langle x'^2\rangle, \langle xx'\rangle, \langle x^2\delta\rangle, \langle x'^2\delta\rangle, \langle xx'\delta\rangle, \ldots$

Summary of Current Limits From Different Focusing Methods

Einzel Lens

$$Q_{max} \cong \frac{3\pi^2}{8}\left(\frac{q\phi_0}{m v_0^2}\right)^2\left(\frac{r_b}{L}\right)^2$$

Solenoids

$$Q_{max} = \left(\frac{\omega_c r_b}{2\gamma\beta c}\right)^2$$

Quadrupole Focusing

$$Q_{max} \cong \frac{\eta\phi_0}{2\pi}\left(\frac{\sin\frac{\eta\pi}{2}}{\frac{\eta\pi}{2}}\right)\left[\frac{B r_b}{[B\rho]}\left[\frac{r_b}{r_p}\right] \quad \frac{2qV_q}{\gamma m v_z^2}\left[\frac{r_b^2}{r_p^2}\right]\right]$$

MAGNETIC

Electric

For Non-Relativistic Beams

$$\lambda_{max} \propto \frac{\phi_0^2}{V} \qquad \lambda_{max} \propto \frac{q}{m} B^2 r_p^2 \qquad \lambda_{max} \propto \begin{cases} B_1 V^{1/2} r_p \\ V_q \end{cases}$$

Note
ϕ_0 = Voltage between Einzel lenses
V_q = Voltage on a quad relative to ground
V = particle energy / q

Solenoids: $\log\{\lambda[Cm^{-1}]\} = -7.0$
Electrostatic quads: $\log\{\lambda[Cm^{-1}]\} = -5.8$
Magnetic quads: $\log\{\lambda[Cm^{-1}]\} = -8.9 + \frac{1}{2}\log\{E_{kin}[eV]\}$
Einzel lenses: $\log\{\lambda[Cm^{-1}]\} = 0.01 - \log\{E_{kin}[eV]\}$



ESTIMATING SPOT SIZE

$$r_x'' + \frac{(\gamma_b \beta_b)'}{\gamma_b \beta_b} r_x + K_x r_x - \frac{2Q}{r_x + r_y} - \frac{\epsilon_x^2}{r_x^3} = 0$$

$$r_y'' + \frac{(\gamma_b \beta_b)'}{\gamma_b \beta_b} r_y + K_y r_y - \frac{2Q}{r_x + r_y} - \frac{\epsilon_y^2}{r_y^3} = 0$$

<u>IN CHAMBER</u>: NO EXTERNAL FOCUSING, NO ACCELERATION AND BEAM IS OFTEN CIRCULAR (BY DESIGN)

$$\Rightarrow \quad K_x = K_y = (\gamma_b \beta_b)' = 0 \qquad \& \quad r_x = r_y = r_b$$

$\Rightarrow$ ENVELOPE EQUATION IS:

$$r_b'' = \frac{Q}{r_b} + \frac{\epsilon^2}{r_b^3}$$



MULTIPLYING BY r_b' & INTEGRATING $\Rightarrow$

$$\frac{r_{bf}'^2}{2} - \frac{r_{bo}'^2}{2} = Q \ln \frac{r_{bf}}{r_{bo}} + \frac{\epsilon^2}{2 r_{bo}^2} - \frac{\epsilon^2}{2 r_{bf}^2}$$

<u>NOW</u> $r_{bo}' \cong \theta$ $\qquad r_{bf}$ = spot radius $\qquad r_{bf} << r_{bo}$

$r_{bf}' = 0 \qquad r_{bo} \cong d\theta$

$$\Rightarrow \boxed{\theta^2 \cong 2Q \ln\left(\frac{\theta d}{r_{bf}}\right) + \frac{\epsilon^2}{r_{bf}^2}}$$

WHEN $Q = 0$

$$r_{spot}^2 = \frac{\epsilon^2}{\theta^2} + r^2_{\text{CHROMATIC ABERRATION}} + \ldots$$

$$r^2_{\text{CHROMATIC}} = \alpha^2 d^2 \left(\frac{\delta p}{p}\right)^2 \theta^2$$

$\alpha \cong 6$ (system dependent)

LONGITUDINAL DYNAMICS SUMMARY

1D VLASOV EQUATION

$$\frac{\partial \tilde{f}}{\partial s} + z' \frac{\partial f}{\partial z} + z'' \frac{\partial \tilde{f}}{\partial z'} = 0$$

$$E_z = \frac{-g}{4\pi\epsilon_0}\frac{\partial \lambda}{\partial z}$$ "g-factor" model

$$z'' = \frac{q E_z}{m v_0^2} \; ; \qquad \frac{\partial^2 \phi}{\partial z^2} + \frac{1}{r}\frac{\partial}{\partial r}\left(r \frac{\partial \phi}{\partial r}\right) = -\frac{\rho}{\epsilon_0}$$

CHILD-LANGMUIR IN 1-D DIODE

LEADS TO FLUID EQUATIONS

$\int$(Vlasov Equation) dz'

$$\frac{\partial \lambda}{\partial s} + \frac{\partial}{\partial z}(\lambda \bar{z}') = 0$$

$$\frac{\partial \bar{z}'}{\partial s} + \bar{z}' \frac{\partial \bar{z}'}{\partial z} + \frac{1}{\lambda}\frac{\partial}{\partial z}\cancel{(\lambda \Delta z'^2)} = \frac{q E_z}{m v_0^2}$$

1D E_z: $\Rightarrow$ CHILD LANGMUIR SOLUTION $\leftarrow$ NON-LINEAR SOLUTION TO FLUID EQUATIONS

2D PIERCE ELECTRODE

TIME DEPENDENT LAMPEL DEFENDACE SOLUTION

g-factor: $\Rightarrow$ SPACE-CHARGE WAVES

$\hookrightarrow$ LONGITUDINAL OR RESISTIVE WALL INSTABILITY (IF $E_1 = z^* I_1$)

$\Rightarrow$ SPACE-CHARGE RAREFACTION WAVES $\leftarrow$ NON-LINEAR SOLUTION TO FLUID EQNS.

$\Rightarrow$ PARABOLIC BUNCH COMPRESSION $\leftarrow$ NON-LINEAR SOLUTION TO FLUID EQUATIONS

$\Rightarrow$ "EAR" FIELDS

VLASOV EQUATION ALSO $\Rightarrow$ ENVELOPE EQUATION $\iint$ VLASOV Equation $dz\,dz'$

$$\frac{d^2 r_z}{ds^2} = \frac{\epsilon_z^2}{r_z^3} + \frac{3}{2}\frac{g q Q_c}{4\pi\epsilon_0 m v^2}\frac{1}{r_z^2} - K(r) r_z$$

KINETIC SOLUTION TO VLASOV EQUATION & SATISFYING RMS ENVELOPE EQUATION IS NEUFER DISTRIBUTION

$$f(z,z') = \frac{3N}{2\pi\epsilon_z}\sqrt{1 - \frac{z^2}{r_z^2} - \frac{r_z^2 (z' - r_z' z/r_z)^2}{\epsilon_z^2}}$$

NORMAL MODES

LONGITUDINAL

SPACE-CHARGE WAVES (FLUID)

$\omega = \pm c_s k$ [IN COMOVING BEAM FRAME]

$$c_s = \sqrt{\frac{q g \lambda_0}{4\pi\epsilon_0 m}}$$ = SPACE CHARGE WAVE SPEED

TRANSVERSE

ENVELOPE MODES

CONTINUOUS FOCUSING (LONG BUNCHES)

BREATHING: $k_B^2 = 2k_{p0}^2 + 2k_p^2$

QUADRUPOLE $k_Q^2 = k_{p0}^2 + 3k_p^2$

(HERE $k_p^2 \equiv k_{p0}^2 - \frac{Q}{r_b^2}$)

(ANALOGOUS MODES IN BUNCHED BEAMS)

STEVE LOOKED AT MODES IN PERIODIC SYSTEMS (& CONTINUOUS FOCUSING

+ KINETIC MODES (GLUCKSTERN MODES)
+ FLUID MODES

INSTABILITIES

1. LONGITUDINAL (RESISTIVE WALL) INSTABILITY
   (FLUID INSTABILITY)

2. ELECTRON – ION INSTABILITY
   (CENTROID INSTABILITY)

STEVE TALKED ABOUT:

3. ENVELOPE INSTABILITIES

STEVE TALKED ABOUT:

4. KINETIC INSTABILITIES
   (DISTRIBUTION FUNCTION DEPENDENT)

5. SINGLE PARTICLE RESONANT INSTABILITIES
   - HALO
   - RING RESONANCES

HALO:

CORE TEST PARTICLE MODEL:

$$x'' = \begin{cases} -\left[k_{\beta 0}^2 - \frac{Q}{r_b^2}\right]x & \text{for } r < r_b \\ -\left[k_{\beta 0}^2 - \frac{Q}{r^2}\right]x & \text{for } r > r_b \end{cases}$$

$$r_b = r_{b0} + \delta r_b \cos(k_\beta s + \phi)$$

Gluckstern's phase-amplitude analysis:

$$x'' + \overbrace{\left[k_{\beta 0}^2 - \frac{Q}{r_{b0}^2}\right]}^{k_\beta^2} x = f(x)$$

Non linear + forcing part

$x = A\sin\psi \quad x' = k_\beta A\cos\psi$ ← PHASE/AMPLITUDE

$\psi = k_\beta s + \alpha$ If $f = 0$ A & ϕ would be constant

$\Rightarrow A' = \frac{1}{k_\beta r_{b0}} f\cos\psi \quad \alpha' = \frac{-1}{k_\beta r_{b0} A} f\sin\psi$

DEFINE RESONANT PHASE $\Psi_r = 2\psi - k_\beta s$

AVERAGE OVER ALL NON-RESONANT FREQUENCIES

$$A_r' = \frac{1}{k_\beta r_{b0}} \int_{-\pi}^{\pi} f\cos\psi \frac{d\psi}{2\pi}; \quad \alpha_r' = \frac{-1}{k_\beta A_r} \int_{-\pi}^{\pi} \frac{d\psi}{2\pi} f\sin\psi$$

$\rightarrow A_r', \Psi_r' \xrightarrow[(\omega = A_r^2)]{} \omega', \Psi_r' \rightarrow H(\omega, \Psi_r) \rightarrow$ CORE RESONANT PARTICLE TRAJECTORY & SEPARATRIX

Numerically determined frequency and amplitude of particle oscillations: linear rf focusing



HI+ Heavy ion	Residual gas molecule	e- electron

Processes:

1. Coulomb collisons (intra-beam)



$$\frac{dT_\perp}{dt} = -\frac{1}{2}\frac{dT_\parallel}{dt} = -\frac{(T_\perp - T_\parallel)}{\tau}$$

$$\tau \sim \nu_c^{-1}$$

2. Coulomb collisions with residual gas



$$\frac{d\sigma_N}{ds} \sim n_g(\quad)$$

3. Charge exchange



4. Stripping



5. Neutralization



CHARGE CHANGING COLLISIONS

Beam loss

+

n_{HI} molecules

6. Gas Ionization



n_g molecules

SUMMARY OF ELECTRON, GAS, PRESSURE, & SCATTERING EFFECTS

1. COULOMB COLLISIONS WITHIN BEAM CAN TRANSFER ENERGY FROM ⊥ TO ∥ AND PROVIDE LOWER LIMIT ON $T_{\parallel}$, HIGHER THAN FROM ACCELERATIVE COOLING.

2. COULOMB INTERACTIONS WITH RESIDUAL GAS NUCLEI PROVIDE A SOURCE OF EMITTANCE GROWTH (BUT NOT IMPORTANT FOR HIGHER MASS AND LONGER RESIDENCE TIMES).

3. PRESSURE INSTABILITY FROM DESORPTION OF RESIDUAL GAS BY STRAYED BEAM IONS HITTING WALL OR BEAM-IONIZED RESIDUAL GAS ATOMS, FORCED TO WALL BY E-field of BEAM. LIMITS CURRENT IN RINGS OR HIGH REP RATE LINAC.

4. ELECTRONS CAN CASCADE AND REACH A "QUASI" EQUILIBRIUM POPULATION OF SIMILAR LINE CHARGE TO THE ION BEAM. ELECTRON-ION TWO STREAM INSTABILITY IS UNSTABLE, AND CAN LEAD TO TRANSVERSE INSTABILITY, SIMILAR TO WHAT IS OBSERVED IN SOME PROTON RINGS.

Inertial Fusion Energy (IFE) power plants of the future will consist of four parts



A power-plant driver would fire about five targets per second to produce as much electricity as today's 1000-megawatt power plant