John Barnard
Steven Lund
USPAS
June 2008

Current limits

A. Axisymmetric
   1. Solenoids
   2. Einzel lens
B. Quadrupolar
   1. Derivation of envelope equations with elliptic symmetry
   2. Current limit using fourier transform method

YESTERDAY, WE DERIVED THE PARAXIAL RAY EQUATION FOR PARTICLES IN AXISYMMETRIC SYSTEMS:

$$\underbrace{r'' + \frac{\gamma'}{\beta^2\gamma} r'}_{\text{INERTIAL}} + \underbrace{\frac{\gamma''}{2\beta^2\gamma}}_{E_r} + \underbrace{\left(\frac{\omega_c}{2\gamma\beta c}\right)^2 r}_{V_\theta B_z \,-\, \text{CENTRIFUGAL}} - \underbrace{\left(\frac{p_\theta}{\gamma\beta m c}\right)^2 \frac{1}{r^3}}_{\text{CENTRIFUGAL}} - \underbrace{\frac{q}{\gamma^3 m v_z^2} \frac{\lambda(r)}{2\pi\epsilon_0 r}}_{\text{SELF-FIELD}} = 0$$

$$\theta' = \frac{p_\theta}{\gamma m r^2 \beta c} - \frac{\omega_c}{2\gamma\beta c}$$ ← CONSTANCY & DEFINITION OF CANONICAL MOMENTUM

ENVELOPE EQUATION FOR AXISYMMETRIC BEAM

$$r_b'' + \frac{\gamma' r_b'}{\beta^2\gamma} + \frac{\gamma''}{2\beta^2\gamma} r_b + \left(\frac{\omega_c}{2\gamma\beta c}\right)^2 r_b - \frac{4\langle p_\theta\rangle^2}{(\gamma m\beta c)^2 r_b^3} - \frac{\varepsilon_r^2}{r_b^3} - \frac{Q}{r_b} = 0$$

$$\varepsilon_r^2 \equiv 4\left(\langle r^2\rangle\langle r'^2\rangle - \langle r r'\rangle^2 + \langle r^2\rangle\langle r^2\theta'^2\rangle - \langle r^2\theta'\rangle^2\right)$$



PERIODIC SOLENOIDS



(FIGURE FROM DAVIDSON & QIN, 2003) p. 55 "PHYSICS OF INTENSE (HIGH-CURRENT) PARTICLE BEAMS IN HIGH ENERGY ACCELERATORS"

Figure 3.2. Schematic of magnet sets producing a periodic focusing solenoidal field with axial periodicity length S. In Fig. 3.2 (a), successive coils are spaced by S and have the same current polarity $+I_1, +I_1, \cdots$. In Fig. 3.2 (b), successive coils are spaced by $S/2$ and have alternating current polarities $+I_1, -I_1, +I_1, \cdots$.

SOLENOIDAL FOCUSING



Let $\gamma' = \gamma'' = 0$

FOR MAXIMUM TRANSPORT $P_\theta = 0$ & $\varepsilon_r^2 = 0$

$$\Rightarrow \quad r_b'' + \left(\frac{\omega_c}{2\gamma\beta c}\right)^2 r_b = \frac{Q}{r_b}$$

FOR A MATCHED BEAM:

$$Q_{max} = \left(\frac{\omega_c}{2\gamma\beta c}\right)^2 r_b^2$$

HEURISTICALLY:



$V_\theta = \omega r$

$$m\omega^2 r + Q m V_z^2 \left(\frac{r}{r_b^2}\right) = q \underbrace{\omega r}_{V_\theta} B_z$$

centrifugal force — SPACE CHARGE FORCE — MAGNETIC FORCE INWARD

$$\Rightarrow \quad \omega^2 + \frac{QV^2}{r_b^2} = \omega\omega_c$$

$\omega\omega_c - \omega^2 =$ MAXIMUM WHEN $\omega = \frac{\omega_c}{2}$

$$\Rightarrow \quad Q_{max} = \left(\frac{\omega_c^2}{4}\right)\left(\frac{r_b^2}{V^2}\right)$$

SOLENOIDAL FOCUSING – CONTINUED

IN REALITY BEAM ACQUIRES V_θ AS BEAM ENTERS SOLENOID:

CONSIDER SIMPLE STEP FUNCTION APPROXIMATION TO SOLENOID FIELD:



LET $B_z = B_0\left[\Theta(z) + \Theta(l_m - z) - 1\right] = \begin{cases} 0 & z < 0 \\ B_0 & 0 < z < l_m \\ 0 & z > l_m \end{cases}$

$$\frac{\partial B_z}{\partial z} = B_0\left[\delta(z) + \delta(l_m - z)\right]$$

HERE $\Theta(z) = \begin{cases} 1 & z > 0 \\ 0 & z < 0 \end{cases}$

AS WE FOUND EARLIER $\nabla \cdot B = 0 \Rightarrow$

$$B_r(r,z) \simeq -\frac{r}{2}\frac{\partial B_z}{\partial z} + \cdots = -\frac{r}{2} B_0\left[\delta(z) + \delta(l_m - z)\right]$$

$$\Delta p_\theta^* = q\int v_z B_r \, dt = \int_{-\infty}^{0+\epsilon} q B_r \, dz = -\frac{r q B_0}{2}$$

$$\Rightarrow v_\theta = r\frac{qB_0}{2m} = \frac{r\omega_c}{2}$$

EINZEL LENS

J. BARNARD

5



FOCUSING OCCURS AT LARGE RADIUS THAN DEFOCUSING

$\Rightarrow$ NET INWARD FORCE

EINZEL LENS – ANALYSIS (DERIVATION FROM ED LEE)

NOW, LET $\omega_c = \langle P_\theta \rangle = \varepsilon_r^2 = 0$

$$\Rightarrow r_b'' + \frac{\gamma'}{\beta^2\gamma} r_b' + \frac{\gamma''}{2\beta^2\gamma} r_b - \frac{Q}{r_b} = 0$$

ALSO ASSUME $\beta \ll 1$, NON-RELATIVISTIC BEAM $\rightarrow \gamma' \simeq \beta\beta'$, $\gamma'' \simeq \beta'^2 + \beta''\beta$

$$r_b'' + \frac{\beta'}{\beta} r_b' + \left[\frac{1}{2}\frac{\beta'^2}{\beta^2} + \frac{1}{2}\frac{\beta''}{\beta}\right] r_b - \frac{Q}{r_b} = 0$$

To eliminate r_b' term try substitution

$$r_b = \left(\frac{\beta_0}{\beta}\right)^{1/2} R$$

$$r_b' = \left(\frac{\beta_0}{\beta}\right)^{1/2} R' - \frac{1}{2}\left(\frac{\beta}{\beta_0}\right)^{-3/2} \frac{R}{\beta_0}\beta'$$

$$r_b'' = \left(\frac{\beta_0}{\beta}\right)^{1/2} R'' - \left(\frac{\beta}{\beta_0}\right)^{-\frac{3}{2}} \frac{R'}{\beta_0}\beta' + \frac{3}{4}\left(\frac{\beta}{\beta_0}\right)^{-\frac{5}{2}} \frac{R}{\beta_0^2}\beta'^2 - \frac{1}{2}\left(\frac{\beta}{\beta_0}\right)^{\frac{3}{2}} \frac{R}{\beta_0}\beta''$$

$$\Rightarrow \left(\frac{\beta_0}{\beta}\right)^{1/2} R'' + \frac{3}{4}\left(\frac{\beta}{\beta_0}\right)^{-\frac{5}{2}} \frac{\beta'^2}{\beta_0^2} R = \frac{Q}{R}\left(\frac{\beta}{\beta_0}\right)^{1/2}$$

$$\Rightarrow \boxed{R'' = \frac{Q}{R}\left(\frac{\beta}{\beta_0}\right) - \frac{3}{4}\left(\frac{\beta'}{\beta}\right)^2 R}$$

EINZEL LENS - CONTINUED

MODEL: LET $\phi = \phi_0 \cos\left(\frac{\pi z}{L}\right)$

$$\frac{1}{2} m v^2 + q\phi = \text{constant}$$

$$\Rightarrow \quad v^2 = v_0^2 + \frac{2q\phi}{m} \cos\left(\frac{\pi z}{L}\right)$$

$$v' = \frac{-q\phi_0}{mv}\left(\frac{\pi}{L}\right)\sin\left(\frac{\pi z}{L}\right)$$

If $\left(\frac{2q\phi_0}{m}\right) << v_0^2$: $\left(\frac{\beta'}{\beta}\right)^2 \cong \left(\frac{q\phi_0}{mv_0^2}\right)^2\left(\frac{\pi}{L}\right)^2 \sin^2\left(\frac{\pi z}{L}\right)$

FOR EQUILIBRIUM LOOK AT D.C. COMPONENT: $\sin^2(x) = \frac{1}{2} - \frac{1}{2}\cos 2x$

$$R'' = 0 \quad \Rightarrow \quad \frac{Q}{\bar{R}} = \frac{3}{4}\overline{\left(\frac{\beta'}{\beta}\right)^2}\,\bar{R}$$

$$R \equiv \left(\frac{\beta}{\beta_0}\right)^{1/2} r_b \quad \Rightarrow \quad \bar{R} = r_b$$

$$\overline{\left(\frac{\beta'}{\beta}\right)^2} = \frac{1}{2}\left(\frac{q\phi_0}{mv_0^2}\right)^2\left(\frac{\pi}{L}\right)^2$$

$$\Rightarrow \quad Q_{max} = \frac{3\pi^2}{8}\left(\frac{q\phi_0}{mv_0^2}\right)^2\left(\frac{r_b}{L}\right)^2$$

CJ. BARNARD
(7.5)

FROM REISER, p.112

$E_x = -E'x$
$E_y = E'y$



$F_x = -qE'x$
$F_y = qE'y$

ELECTROSTATIC QUADS

Figure 3.15. Electrodes and force lines in an electrostatic quadrupole.

$B_x = B'y$
$B_y = B'x$

$F_x = -qV_z B'x$
$F_y = qV_z B'y$



MAGNETIC QUADS

EQUATION OF MOTION

RETURN TO X, Y COORDINATES

$$x'' + \frac{1}{\gamma v_z}\frac{d}{ds}(\gamma v_z)x' = \frac{-q}{\gamma^3 m v_z^2}\frac{\partial \varphi}{\partial x} \pm \begin{cases} \frac{qB'}{\gamma m v_z^2}x & \text{for magnetic quads} \\ \frac{qE'}{\gamma m v_z^2}x & \text{for electric quads} \end{cases}$$

Let $\frac{\gamma m v_z}{q} = \frac{p}{q} \equiv [B\rho] \equiv$ RIGIDITY

$$y'' + \frac{1}{\gamma v_z}\frac{d}{ds}(\gamma v_z)y' = \frac{-q}{\gamma^3 m v_z^2}\frac{\partial \varphi}{\partial y} \mp \begin{cases} \frac{B'}{[B\rho]}y & \text{magnetic} \\ \frac{qE'}{\gamma m v_z^2}y & \text{electric} \end{cases}$$

ENVELOPE EQUATION

$$r_x^2 = 4\langle x^2\rangle; \quad r_y^2 = 4\langle y^2\rangle$$

$$r_x' = \frac{4\langle xx'\rangle}{r_x}$$

$$r_x'' = \frac{4\langle xx''\rangle}{r_x} + \frac{\varepsilon_x^2}{r_x^3}; \qquad \varepsilon_x^2 = 16\left(\langle x^2\rangle\langle x'^2\rangle - \langle xx'\rangle^2\right)$$

$$r_y'' = \frac{4\langle yy''\rangle}{r_y} + \frac{\varepsilon_y^2}{r_y^3} \qquad \varepsilon_y^2 = 16\left(\langle y^2\rangle\langle y'^2\rangle - \langle yy'\rangle^2\right)$$

for magnetic focusing:

$$r_x'' + \frac{1}{\gamma v_z}\frac{d}{ds}(\gamma v_z)r_x' + \frac{4q}{\gamma^3 m v_z^2}\frac{\langle x\frac{\partial\varphi}{\partial x}\rangle}{r_x} \pm \frac{B'}{[B\rho]}r_x - \frac{\varepsilon_x^2}{r_x^3} = 0$$

$$r_y'' + \frac{1}{\gamma v_z}\frac{d}{ds}(\gamma v_z)r_y' + \frac{4q}{\gamma^3 m v_z^2}\frac{\langle y\frac{\partial\varphi}{\partial y}\rangle}{r_y} \pm \frac{B'}{[B\rho]}r_x - \frac{\varepsilon_y^2}{r_y^3} = 0$$

(for electric focusing $\frac{B'}{[B\rho]} \to \frac{qE'}{\gamma m v_z^2}$)

SPACE CHARGE TERM WITH ELLIPTICAL SYMMETRY

FY: ELLIPTICAL SYMMETRY: $\rho = \rho\left(\frac{x^2}{r_x^2} + \frac{y^2}{r_y^2}\right)$

CAN BE SHOWN THAT $\left\langle x \frac{\partial \phi}{\partial x} \right\rangle = \frac{-\lambda}{4\pi\varepsilon_0} \frac{r_x}{r_x + r_y}$

$$\left\langle y \frac{\partial \phi}{\partial y} \right\rangle = \frac{-\lambda}{4\pi\varepsilon_0} \frac{r_y}{r_x + r_y}$$

DEFINING $Q = \frac{2q\lambda}{4\pi\varepsilon_0 \gamma^3 m v_z^2}$

$$\boxed{\begin{aligned} r_x'' + \frac{1}{\gamma v_z}\frac{d}{ds}(\gamma v_z) r_x' - \frac{2Q}{r_x + r_y} \mp \frac{B'}{[B\rho]} r_x - \frac{\varepsilon_x^2}{r_x^3} &= 0 \\ r_y'' + \frac{1}{\gamma v_z}\frac{d}{ds}(\gamma v_z) r_y' - \frac{2Q}{r_x + r_y} \pm \frac{B'}{[B\rho]} r_y - \frac{\varepsilon_y^2}{r_y^3} &= 0 \end{aligned}}$$

(for Electric Focusing $\frac{B'}{[B\rho]} \to \frac{qE'}{\gamma m v_z^2}$).

SPACE CHARGE TERM WITH ELLIPTICAL SYMMETRY II

ELLIPTICAL SYMMETRY: $\rho = \rho\left(\frac{x^2}{r_x^2} + \frac{y^2}{r_y^2}\right)$

CAN BE SHOWN THAT (Sacherer, 1971)

$$\left\langle x \frac{\partial \varphi}{\partial x} \right\rangle = \frac{-\lambda}{4\pi\varepsilon_0} \frac{r_x}{r_x + r_y}$$

&

$$\left\langle y \frac{\partial \varphi}{\partial y} \right\rangle = \frac{-\lambda}{4\pi\varepsilon_0} \frac{r_y}{r_x + r_y}$$

OUTLINE OF PROOF: (from R. Ryne)

Let $\chi = \frac{x^2}{r_x^2+s} + \frac{y^2}{r_y^2+s}$

DEFINE $\eta(\chi)$ such that $\rho(x,y) = \left.\frac{d\eta(\chi)}{d\chi}\right|_{s=0} = \hat{\rho}(\chi)\big|_{s=0}$

So $\rho = \hat{\rho}\left(\frac{x^2}{r_x^2} + \frac{y^2}{r_y^2}\right) = \hat{\rho}(\chi)\big|_{s=0}$

DEFINE $\Psi(x,y) = \frac{-r_x r_y}{4\varepsilon_0} \int_0^\infty \frac{\eta(\chi)\, ds}{\sqrt{r_x^2+s}\,\sqrt{r_y^2+s}}$

It follows that $\frac{\partial^2\Psi}{\partial x^2} + \frac{\partial^2\Psi}{\partial y^2} = -\frac{\rho}{\varepsilon_0}$ AND SO IS A SOLUTION OF POISSON'S EQUATION (since $\Psi \to 0$ as $x,y \to \infty$)

WHAT IS $\left\langle x \frac{\partial\Psi}{\partial x} \right\rangle$?

$$\left\langle x \frac{\partial\Psi}{\partial x} \right\rangle = \frac{-r_x r_y}{4\lambda\varepsilon_0} \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} dy\, x\, \rho(x,y) \int_0^\infty \frac{\eta' \frac{\partial\chi}{\partial x}\, ds}{\sqrt{r_x^2+s}\,\sqrt{r_y^2+s}}$$

where $\lambda = \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} dy\, \rho(x,y)$

So $$\left\langle x \frac{\partial \psi}{\partial x} \right\rangle = \frac{-2 r_x r_y}{4 \lambda \varepsilon_0} \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} dy \; x^2 \hat{\rho}\left(\frac{x^2}{r_x^2} + \frac{y^2}{r_y^2}\right) \int_0^{\infty} \frac{\hat{\rho}\left(\frac{x^2}{r_x^2+s} + \frac{y^2}{r_y^2+s}\right) ds}{(r_x^2+s)^{3/2}(r_y^2+s)^{1/2}}$$

Let $$r\cos\theta = \frac{x}{\sqrt{r_x^2+s}} \qquad r\sin\theta = \frac{y}{\sqrt{r_y^2+s}}$$

$$\det J = \sqrt{r_x^2+s}\,\sqrt{r_y^2+s}\; r$$ where J is the Jacobian $dx\,dy = \det J \cdot dr\,d\theta$

$$\Rightarrow \left\langle x \frac{\partial \psi}{\partial x} \right\rangle = \frac{-r_x r_y}{\lambda 2 \varepsilon_0} \int_0^{\infty} dr \int_0^{2\pi} d\theta \int_0^{\infty} ds \; r^3 \hat{\rho}(r^2)\, \rho\left(\frac{r_x^2+s}{r_x^2} r^2\cos^2\theta + \frac{r_y^2+s}{r_y^2} r^2 \sin^2\theta\right) \cdot \cos^2\theta$$

Let $$r'^2 = \frac{r_x^2+s}{r_x^2} r^2\cos^2\theta + \frac{r_y^2+s}{r_y^2} r^2 \sin^2\theta$$
$$= r^2\left[1 + s\left(\frac{\cos^2\theta}{r_x^2} + \frac{\sin^2\theta}{r_y^2}\right)\right]$$

with r fixed $$2r'dr' = r^2\left(\frac{\cos^2\theta}{r_x^2} + \frac{\sin^2\theta}{r_y^2}\right) \cdot ds$$

$$\Rightarrow \left\langle x \frac{\partial \psi}{\partial x} \right\rangle = \frac{-r_x r_y}{2\lambda \varepsilon_0} \int_0^{\infty} dr \int_0^{2\pi} d\theta \int_r^{\infty} dr' \frac{2r'dr'\, r^3\, \hat{\rho}(r^2)\, \hat{\rho}(r'^2)}{r^2\left(\frac{\cos^2\theta}{r_x^2} + \frac{\sin^2\theta}{r_y^2}\right)} \cos^2\theta$$

$$\int_0^{2\pi} \frac{\cos^2\theta\, d\theta}{\frac{\cos^2\theta}{r_x^2} + \frac{\sin^2\theta}{r_y^2}} = \frac{2\pi r_x^2 r_y}{r_x + r_y}$$

$$\Rightarrow \left\langle x \frac{\partial \psi}{\partial x} \right\rangle = \frac{-r_x^3 r_y^2}{\lambda 2\pi \varepsilon_0 (r_x + r_y)} \int_0^{\infty} dr\, 2\pi r\, \hat{\rho}(r^2) \int_r^{\infty} dr'\, 2\pi r'\, \hat{\rho}(r'^2)$$

Recall $$\lambda = \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} dx\,dy\, \rho(x,y) = \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} dx\,dy\, \hat{\rho}\left(\frac{x^2}{r_x^2} + \frac{y^2}{r_y^2}\right)$$

Let $$\frac{x}{r_x} = r\cos\theta \qquad \frac{y}{r_y} = r\sin\theta \qquad \det J = r_x r_y r$$
$$\Rightarrow \lambda = \int_0^{\infty} dr \int_0^{2\pi} d\theta\, \rho(r^2)\, r_x r_y r = 2\pi r_x r_y \int_0^{\infty} dr\, r\, \rho(r^2)$$

Now $\displaystyle\int_0^\infty dr\, r\, \rho(r^2) \int_0^r dr'\, r'\, \rho(r'^2) = \frac{1}{2} \overbrace{\int_0^\infty dr\, r\, \rho(r^2)}^{\lambda/2\pi r_x r_y} \overbrace{\int_0^\infty dr'\, r'\, \rho(r'^2)}^{\lambda/2\pi r_x r_y}$



(by symmetry & consideration of diagram at left.)

$$\Rightarrow \quad \left\langle x \frac{\partial \psi}{\partial x} \right\rangle = \frac{-\lambda}{4\pi\epsilon_0} \frac{r_x}{r_x + r_y}$$

National® Brand 42-182 100 SHEETS Made in U.S.A.

CURRENT LIMIT FOR QUADRUPOLES



$$k = \begin{cases} \dfrac{B_0'}{[B\rho]} & \text{MAGNETIC} \\ \dfrac{qE_0'}{\gamma m v_z^2} & \text{ELECTRIC} \end{cases}$$

$$r_x'' + k f(s) r_x - \frac{2Q}{r_x + r_y} = 0$$

(NOTE WE HAVE SET $\epsilon = 0$).

$$r_y'' - k f(s) r_y - \frac{2Q}{r_x + r_y} = 0$$

$$f(s) = \begin{cases} 1 & 0 < s < \eta L/2 \\ -1 & L - \frac{\eta L}{2} < s < L + \eta L/2 \\ 1 & 2L - \eta L/2 < s < 2L \\ 0 & \text{otherwise} \end{cases}$$

$$\frac{1}{L}\int_0^{2L} f(s) \cos\left(\frac{\pi s}{L}\right) ds = \left(\frac{4}{\pi}\right)\sin\left(\frac{\pi\eta}{2}\right)$$ = fourier amplitude at fundamental lattice frequency

Let $r_x = r_b\left(1 + \delta\cos\left(\frac{\pi s}{L}\right)\right)$

$r_y = r_b\left(1 - \delta\cos\left(\frac{\pi s}{L}\right)\right)$

Let $k f(s) \rightarrow k\left(\frac{4}{\pi}\right)\sin\left(\frac{\eta\pi}{2}\right) \cos\left(\frac{\pi s}{L}\right)$

COLLECTING "FAST" TERMS AND "SLOW" TERMS

$$\left[\frac{-\pi^2}{L^2} r_b \delta + k r_b \left(\frac{4\sin\left(\frac{\eta\pi}{2}\right)}{\pi}\right)\right]\cos\left(\frac{\pi s}{L}\right) = 0 \qquad \text{(fast)}$$

$$\delta k r_b \left(\frac{2\sin\left(\frac{\eta\pi}{2}\right)}{\pi}\right) = \frac{Q}{r_b} \qquad \text{(slow)}$$

Fast $\Rightarrow$ $\delta = \dfrac{4kL^2}{\pi^3}\sin\left(\dfrac{\eta\pi}{2}\right)$ & $Q_{max} \cong \dfrac{2\eta^2 k^2 L^2}{\pi^2}\left(\dfrac{\sin\left(\frac{\eta\pi}{2}\right)}{\left(\frac{\eta\pi}{2}\right)}\right)^2 r_b^2$

CONTINUOUS FOCUSING

$$r_x'' = -k_{\beta 0}^2 r_x + \frac{2Q}{r_x + r_y} - \frac{\epsilon^2}{r_x^2}$$

$$r_y'' = -k_{\beta 0}^2 r_y + \frac{2Q}{r_x + r_y} - \frac{\epsilon^2}{r_y^2}$$

CURRENT LIMIT BALANCES PERVEANCE & EXTERNAL FOCUSING $(r_x = r_y = r_b)$:

$$k_{\beta 0}^2 r_b = \frac{Q_{max}}{r_b}$$

Effective $k_{\beta 0}^2$ FOR QUADRUPOLES FOUND FROM DOMINANT FOURIER COMPONENT

$$k_{\beta 0}^2 = \frac{2\eta^2 k^2 L^2}{\pi^2} \left(\frac{\sin\left(\frac{\eta\pi}{2}\right)}{\frac{\eta\pi}{2}} \right)^2 \qquad \text{where } k = \frac{B'}{[B\rho]}$$

FOR CONTINUOUS FOCUSING: $k_{\beta 0}^2 = \frac{\sigma_0^2}{4L^2}$

ELIMINATING L:

$$Q_{max} = \frac{\eta k \sigma_0}{\sqrt{2}\pi} \left(\frac{\sin\frac{\eta\pi}{2}}{\frac{\eta\pi}{2}} \right) r_b^2 \longleftarrow$$

PERVEANCE LIMIT FOR FODO QUADRUPOLES

Envelope instabilities set upper limit on "single particle" phase advance σ_0

Experimental data (Tiefenback, 1986) from LBL's Single Beam Transport Experiment (SBTE)

$\sigma_0 < 85^0$

Experimental limits on beam stability in terms of σ and σ_0



SEE LUND & CHAWLA 2006, NIMPR-A, FOR HIGHER ORDER PARTICLE-LATTICE RESONANCES WHICH CLARIFIES $\sigma_0 = 85^0$ LIMIT

SEE STRUCKMEIER & REISER, PARTICLE ACCELERATORS 14, 227, (1984) & LUND & BUKH, PRSTAB, 7 024801 (2004)

J. BARNARD

QUADRUPOLE CURRENT LIMIT — CONTINUED

$$Q_{max} \cong \frac{\eta k Q_0}{\sqrt{2\pi}} \left(\frac{\sin \frac{\eta \pi}{2}}{\frac{\eta \pi}{2}} \right) r_b^2$$

here $k = \begin{cases} \dfrac{dB/dx}{[B_\rho]} \sim \dfrac{B}{[B_\rho]\, r_p} & \text{(MAGNETIC QUAD FODO)} \\ \dfrac{q\, dE/dx}{\gamma m v_z^2} \sim \dfrac{2 q V_q}{\gamma m v_z^2 r_p^2} & \text{where } V_q = \frac{1}{2}\frac{dE}{dx} r_p^2 \quad \text{(ELECTRIC QUAD FODO)} \end{cases}$

So

$$Q_{max} \cong \frac{\eta Q_0}{\sqrt{2\pi}} \left(\frac{\sin \frac{\eta \pi}{2}}{\frac{\eta \pi}{2}} \right) \begin{cases} \dfrac{B\, r_b}{[B_\rho]} \left[\dfrac{r_b}{r_p} \right] & \text{(MAGNETIC QUAD)} \\ \dfrac{2 q V_q}{\gamma m v_z^2} \left[\dfrac{r_b^2}{r_p^2} \right] & \text{(ELECTRIC QUAD)} \end{cases}$$

Summary of Current Limits From Different Focusing Methods

Einzel Lens

$$Q_{max} \cong \frac{3\pi^2}{8}\left(\frac{q\phi_0}{m v_0^2}\right)^2\left(\frac{r_b}{L}\right)^2$$

Solenoids

$$Q_{max} = \left(\frac{\omega_c r_b}{2\gamma\beta c}\right)^2$$

Quadrupole Focusing

$$Q_{max} \cong \frac{\eta\phi_0}{\sqrt{2}\pi}\left(\frac{\sin\frac{\eta\pi}{2}}{\frac{\eta\pi}{2}}\right)\left[\frac{B r_b}{[B\rho]}\left[\frac{r_b}{r_p}\right] \quad \frac{2qV_q}{\gamma m v_z^2}\left[\frac{r_b^2}{r_p^2}\right]\right]$$

MAGNETIC

Electric

For Non-Relativistic Beams

$$\lambda_{max} \propto \frac{\phi_0^2}{V} \qquad \lambda_{max} \propto \frac{q}{m} B^2 r_p^2 \qquad \lambda_{max} \propto \begin{cases} B_1 V^{1/2} r_p \\ V_q \end{cases}$$

Note
ϕ_0 = Voltage between Einzel lenses
V_q = Voltage on a quad relative to ground
V = particle energy / q